

## CLASSIFICATION PLOTS

<i>Menu item</i>	<i>Module name</i>	<i>Scope<sup>1</sup></i>	<i>Plot description</i>	<i>Details (reference)</i>
AFM (Irvine + Baragar 1971)	AFM	G	(Na <sub>2</sub> O+K <sub>2</sub> O) – FeOt – MgO ternary	AFM plot that serves to discriminate between calc-alkaline and tholeiitic subalkaline series (Irvine & Baragar, 1971).
SiO <sub>2</sub> - FeOt/MgO (Miyashiro 1974)	Miyashiro	G	SiO <sub>2</sub> vs. FeOt /MgO binary	Diagram of Miyashiro (1974) distinguishing between tholeiitic and calc-alkaline igneous rocks.
SiO <sub>2</sub> - K <sub>2</sub> O (Peccerillo + Taylor 1976)	PeceTaylor	G	SiO <sub>2</sub> vs. K <sub>2</sub> O binary	Diagram proposed by Peccerillo & Taylor (1976) to distinguish various series of tholeiitic, calc-alkaline and shoshonitic rocks.
Co - Th (Hastie et al. 2007)	Hastie	G	Co vs. Th	Replacement for the previous plot of Peccerillo & Taylor (1976) using less mobile elements, designed by Hastie <i>et al.</i> (2007).
Molar Na <sub>2</sub> O – Al <sub>2</sub> O <sub>3</sub> – K <sub>2</sub> O plot	NaAlK	G	Na <sub>2</sub> O – Al <sub>2</sub> O <sub>3</sub> – K <sub>2</sub> O ternary	Diagram to distinguish meta-/peraluminous from peralkaline rocks as well as potassic, sodic and ultrapotassic suites.
A/CNK - A/NK (Shand 1943)	Shand	G	Al <sub>2</sub> O <sub>3</sub> /(CaO+Na <sub>2</sub> O+K <sub>2</sub> O) vs. Al <sub>2</sub> O <sub>3</sub> /(Na <sub>2</sub> O+K <sub>2</sub> O) (mol. %)	Classic A/CNK vs A/NK plot of Shand (1943) discriminating metaluminous, peraluminous and peralkaline compositions.
TAS (Le Bas et al. 1986)	TAS	V	SiO <sub>2</sub> vs. (Na <sub>2</sub> O + K <sub>2</sub> O) binary	The principal variation of the TAS diagram, as proposed by Le Bas <i>et al.</i> (1986) and codified by Le Maitre (1989). Dividing line between alkaline and subalkaline series is that of Irvine & Baragar (1971).
TAS (Cox et al. 1979)	CoxVolc CoxPlut	V P	SiO <sub>2</sub> vs. (Na <sub>2</sub> O + K <sub>2</sub> O) binary	Variation of the TAS diagram proposed by Cox <i>et al.</i> (1979) and adopted by Wilson (1989) for plutonic rocks.
TAS (Middlemost 1994)	TASMiddlemostVolc TASMiddlemostPlut	V P	SiO <sub>2</sub> vs. (Na <sub>2</sub> O + K <sub>2</sub> O) binary	Variation of the TAS diagram proposed by Middlemost (1994).
Jensen (1976)	Jensen	V	Al – (Fe <sup>t</sup> + Ti) – Mg ternary	Ternary plot of Jensen (1976).
R1-R2 (De la Roche et al. 1980)	LarocheVolc LarochePlut	V P	R <sub>1</sub> –R <sub>2</sub> binary (in millications).	Multicationic classification plot of De La Roche <i>et al.</i> (1980) (R <sub>1</sub> : 4Si - 11(Na + K) – 2(Fe + Ti); R <sub>2</sub> : 6Ca + 2Mg + Al).
Nb/Y - Zr/TiO <sub>2</sub> (Winchester + Floyd 1977)  Zr/TiO <sub>2</sub> - SiO <sub>2</sub> (Winchester + Floyd 1977)	WinFloyd1  WinFloyd2	V	log Nb/Y vs. log Zr/TiO <sub>2</sub>  log Zr/TiO <sub>2</sub> vs. SiO <sub>2</sub> binary	Diagrams proposed by Winchester & Floyd (1977) for classification of volcanic rocks using incompatible element ratios.
Nb/Y - Zr/Ti plot (modified by Pearce 1996)	Pearce1996	V	log Nb/Y vs. log Zr/Ti	The log Nb/Y vs. log Zr/TiO <sub>2</sub> plot of Winchester & Floyd (1977) modified by Pearce (1996).
QAPF diagram (Streckeisen 1978)	QAPFVolc	V	QAPF – modal compositions	Modal QAPF diagram of Streckeisen (1978)
QAPF diagram (Streckeisen 1974)	QAPFPlut	P	QAPF – modal compositions	Modal QAPF diagram of Streckeisen (1974)

<sup>1</sup>Scope: G: general diagram, V: designed for volcanic rocks, P: designed for plutonic rocks

## CLASSIFICATION PLOTS (CONTD.)

<i>Menu item</i>	<i>Module name</i>	<i>Scope<sup>1</sup></i>	<i>Plot description</i>	<i>Details (reference)</i>
Feldspar triangle (O'Connor 1965)	OConnorVolc OConnorPlut	V P	Ternary plot Ab-An-Or	Classification diagram after O'Connor (1965) for silica-rich rocks (quartz > 10 %). It is based on CIPW-normative (volcanic, plutonic rocks) or modal (plutonic rocks) contents of albite, anorthite and K-feldspar.
P-Q (Debon + Le Fort 1983)	DebonPQ	P	P–Q binary (in millications)	Nomenclature diagram of Debon & Le Fort (1983). Its coordinates correspond to proportions of Kfs and Pl to Qtz (P: K - (Na + Ca), Q: Si/3 - (K + Na + 2Ca/3)).
B-A (Debon + Le Fort 1983)	DebonBA	P	B–A binary (in millications)	The B–A diagram (Debon & Le Fort 1983) defines six sectors (I - VI), reflecting alumina balance of samples (B: Fe + Mg + Ti, A: Al - (K + Na + 2Ca)).
B-A plot (modified by Villaseca et al. 1998)	Villaseca	P	B–A binary (in millications)	The B–A diagram (Debon & Le Fort 1983) with fields of various types of peralkaline rocks as outlined by Villaseca <i>et al.</i> (1998)
Middlemost (1985)	MiddlemostPlut	P	SiO <sub>2</sub> vs. (Na <sub>2</sub> O + K <sub>2</sub> O) binary	Classification diagram of Middlemost (1985) for plutonic rocks.
(or+ab)-2Q-4an diagram (Enrique 2018)	Enrique	P	Ternary plot (Or + Ab)–2Q–4An	CIPW-normative ternary plot of Enrique (2018)
Q-ANOR diagram (Streckeisen + Le Maitre 1979)	QANOR	P	Q' = 100Qz/(Qz + Or + Ab + An) vs. ANOR = 100An/(Or + An) binary b	Q'–ANOR classification diagram of Streckeisen & Le Maitre (1979) based on improved granitoid mesonorm of Mielke & Winkler (1979).

<sup>1</sup>Scope: G: general diagram, V: volcanic rocks, P: plutonic rocks

## GEOTECTONIC PLOTS

<i>Menu item</i>	<i>Module name</i>	<i>Scope<sup>1</sup></i>	<i>Plot description</i>	<i>Details (reference)</i>
Batchelor + Bowden (1985)	Batchelor	Gr	R <sub>1</sub> –R <sub>2</sub> binary (in millications)	R <sub>1</sub> –R <sub>2</sub> diagram (De La Roche <i>et al.</i> , 1980) with geotectonic implications after Batchelor & Bowden (1985). (R <sub>1</sub> : 4Si – 11(Na + K) – 2(Fe + Ti); R <sub>2</sub> : 6Ca + 2Mg + Al).
Source diagram (Laurent et al. 2014)	LaurentSource	G	Ternary plot Al <sub>2</sub> O <sub>3</sub> /(FeOt+MgO), 3CaO and 5K <sub>2</sub> O/Na <sub>2</sub> O (all in wt. %)	Diagram to decipher possible sources of granitic magmas ( Laurent <i>et al.</i> , 2014).
Maniar + Piccoli (1989)	Maniar	Gr	binary plots SiO <sub>2</sub> vs. K <sub>2</sub> O, Al <sub>2</sub> O <sub>3</sub> , and FeOt/(FeOt+MgO); MgO vs. FeOt; CaO vs. FeOt+MgO; A/CNK vs. A/NK	Major-element based geotectonic classification of granitoids (Maniar & Piccoli, 1989).
Frost et al. (2001)	Frost	Gr	binary plots SiO <sub>2</sub> vs. FeO(t)/(FeO(t)+MgO) wt. %; SiO <sub>2</sub> vs. MALI (Na <sub>2</sub> O+K <sub>2</sub> O-CaO) ASI vs. A/NK	Major-element based classification of granitoids (Frost <i>et al.</i> , 2001).
Frost + Frost (2008)	Frost_2008	Gr	binary plots SiO <sub>2</sub> vs. FeOt/(FeOt+MgO) wt. %; SiO <sub>2</sub> vs. MALI (Na <sub>2</sub> O+K <sub>2</sub> O-CaO) SiO <sub>2</sub> vs. ASI, FSSI vs. AI	Major-element based classification of granitoids (Frost & Frost 2008)
A type granitoids (Whalen et al. 1987)	Whalen	Gr	binary plots Zr+Nb+Ce+Y vs. FeOt/MgO (Na <sub>2</sub> O+K <sub>2</sub> O)/CaO 10000×Ga/Al vs. (Na <sub>2</sub> O+K <sub>2</sub> O) (Na <sub>2</sub> O+K <sub>2</sub> O)/CaO, K <sub>2</sub> O/MgO, FeOt/MgO, Zr, Nb, Ce, Y, Zn, Agpaitic index	Binary plots serving for distinction of A-type granitoid rocks after Whalen <i>et al.</i> (1987).
Pearce et al. (1984)	Pearce_granite	Gr	log(Y+Nb) vs. log Rb, log Y vs. log Nb, log(Ta+Yb) vs. log Rb, log Yb vs. log Ta	Trace-element based geotectonic classification of granitoids by Pearce <i>et al.</i> (1984).
Harris et al. (1986)	Harris	Gr	ternary plot Hf - Rb/30 - Ta×3	The diagram distinguishes among four types of collisional granites.
Sylvester (1989)	Sylvester	Gr	(Al <sub>2</sub> O <sub>3</sub> +CaO)/(FeOt+Na <sub>2</sub> O+K <sub>2</sub> O) vs. 100×(MgO+FeOt+TiO <sub>2</sub> )/SiO <sub>2</sub>	Diagram proposed by Sylvester (1989) to distinguish the alkaline collision- related alkaline granites.
Schandl + Gorton (2002)	Schandl	Gr	log Ta/Yb vs. log Th/Ta Ta vs. Th, Ta/Hf vs. Th/Hf Yb vs. Th/Ta	Discrimination of geotectonic environment of felsic volcanic rocks (rhyolites), proposed by Schandl & Gorton (2002). It is based on combination of four presumably little immobile trace elements (Ta, Yb, Th, Hf).

## GEOTECTONIC PLOTS (CONTD.)

<i>Menu item</i>	<i>Module name</i>	<i>Scope<sup>1</sup></i>	<i>Plot description</i>	<i>Details (reference)</i>
YbN vs. LaN/YbN (Martin 1986) TTG/adakite	LaYb	Gr	binary plot Yb <sub>N</sub> vs. La <sub>N</sub> /Yb <sub>N</sub>	Diagram discriminating between adakitic (or TTG) and "ordinary" calc-alkaline rocks (Martin, 1986).
NbN - ThN (Saccani 2015)	Saccani	B	binary plot Nb <sub>N</sub> -Th <sub>N</sub> (NMORB-normalized values)	Detailed discrimination of geotectonic environment of basaltic rocks, proposed by Saccani (2015).
Verma et al. (2006) based on major elements	Verma	B	suite of five diagrams based on log-transformed concentration ratios of major-element oxides	Discrimination of geotectonic environment of ultrabasic and basic rocks (SiO <sub>2</sub> < 52 wt. %), proposed by Verma <i>et al.</i> (2006).
Agrawal et al. (2008), La, Sm, Yb, Nb, Th based	Agrawal	B	suite of five diagrams based on log-transformed concentration ratios of La, Sm, Yb, Nb and Th	Discrimination of geotectonic environment of ultrabasic and basic rocks, proposed by Agrawal <i>et al.</i> (2008). It is based on log-transformed concentration ratios of five trace elements (La, Sm, Yb, Nb, and Th), i.e., using four ratios ln(La/Th), ln(Sm/Th), ln(Yb/Th), and ln(Nb/Th).
Meschede (1986) Zr/4-2Nb-Y	Meschede	B	Zr/4 – 2Nb – Y ternary	(Meschede, 1986)
Mullen (1983) 10MnO-TiO <sub>2</sub> -10P <sub>2</sub> O <sub>5</sub>	Mullen	B	10 MnO – TiO <sub>2</sub> – 10 P <sub>2</sub> O <sub>5</sub>	(Mullen, 1983)
Pearce + Cann (1973)	Pearce_and_Cann	B	Zr – Ti/100 – 3 Y ternary, Zr – Ti/100 – Sr/2 ternary, log Zr – log Ti binary	(Pearce & Cann, 1973)
Pearce + Norry (1979)	Pearce_and_Norry	B	log Zr vs. log Zr/Y	(Pearce & Norry, 1979)
Pearce et al. (1977) MgO-FeOt-Al <sub>2</sub> O <sub>3</sub>	Pearce_et_al_1977	B	MgO – FeOt – Al <sub>2</sub> O <sub>3</sub> ternary	(Pearce <i>et al.</i> , 1977)
Pearce (1982)	Pearce_1982	B	log Zr vs. log. Ti	(Pearce, 1982)
Shervais (1982)	Shervais	B	log Ti/1000 vs. log V	(Shervais, 1982)
Wood (1980)	Wood	B	Th – Hf/3 – Ta Th – Hf/3 – Nb/16 Th – Zr/117 – Nb/16	(Wood, 1980)
Hollocher et al. (2012) La/Yb vs. Nb/La La/Yb vs. Th/Nb	Hollocher1 Hollocher2	B	La/Yb – Nb/La La/Yb – Th/Nb	(Hollocher et al., 2012)
Pearce (2008) Nb/Yb - Th/Yb	PearceNbThYb	U	Nb/Yb – Th/Yb	(Pearce, 2008)
Pearce (2008) Nb/Yb - TiO <sub>2</sub> /Yb	Nb/Yb - TiO <sub>2</sub> /Yb	U	Nb/Yb – TiO <sub>2</sub> /Yb	(Pearce, 2008)

## GEOTECTONIC PLOTS (CONTD.)

<i>Menu item</i>	<i>Module name</i>	<i>Scope<sup>1</sup></i>	<i>Plot description</i>	<i>Details (reference)</i>
Cabanis + Lecolle (1989) La/10-Y/15-Nb/8	Cabanis	U	La/10 – Y/15-Nb/8	(Cabanis & Lecolle, 1989)
Ross + Bedard (2009) Zr/Y-Th/Yb	Ross	U	Zr/Y – Th/Yb	(Ross & Bédard, 2009)
Müller et al. (1992) Potassic rocks binary	MullerKbinary	U		(Müller <i>et al.</i> , 1992)
Müller et al. (1992) Potassic rocks ternary	MullerKternary	U		(Müller <i>et al.</i> , 1992)
Ohta + Arai (2007) FMW weathering index	OhtaArai	U	Ternary plot F – M – W, see help	F-M-W diagram (Ohta & Arai, 2007) for chemical weathering in rocks.

## USER-DEFINED PLOTS

<i>Menu item</i>	<i>Module name</i>	<i>Scope<sup>1</sup></i>	<i>Plot description</i>	<i>Details (reference)</i>
Ta/Yb - K <sub>2</sub> O/Yb (Pearce 1982)	PearceDestructive1	U	Ta/Yb vs. K <sub>2</sub> O/Yb binary diagram	(Pearce, 1982)
Ta/Yb - Th/Yb (Pearce 1982)	PearceDestructive2	U	Ta/Yb vs. Th/Yb binary diagram	(Pearce, 1982)
Nb/Y - Ti/Y (Pearce 1982)	PearceDestructive3	U	Nb/Y vs. Ti/Y binary diagram	(Pearce, 1982)
Al <sub>2</sub> O <sub>3</sub> /SiO <sub>2</sub> - MgO/SiO <sub>2</sub> (Paulick et al. 2006)	Paulick	UB	Al <sub>2</sub> O <sub>3</sub> /SiO <sub>2</sub> vs. MgO/SiO <sub>2</sub> binary plot for mantle peridotites	(Paulick <i>et al.</i> , 2006)
K/(Na+K)-B (Debon + Le Fort 1988)	DebonKNaB	U	K/(Na+K) vs. B binary (in millications)	The K/(Na+K) vs. B (maficity) diagram of Debon & Le Fort (1988) for aluminous magmatic suites. It defines three associations based on the balance of the two alkalis, namely potassic, sodi-potassic and sodic.
Q-B (Debon + Le Fort 1988)	DebonQB	U	Q vs. B binary (in millications)	The Q (cation proportion of quartz) vs. B (maficity) diagram of Debon & Le Fort (1983) for aluminous associations. It defines three associations, reflecting maficity of samples: leucocratic, subleucocratic and mesocratic.
B-Mg.no (Debon + Le Fort 1988)	DebonBMgNo	U	B vs. Mg# binary (in millications)	Diagram of Debon & Le Fort (1983) of B (maficity) vs. Mg/(Fe + Mg) to distinguish magnesian and ferriferrous associations.
B-Q-F (Debon + Le Fort 1988)	DebonBQF	U	B–Q–F terary (in millications)	Diagram of Debon & Le Fort (1983) expresses balance of three main groups of rock-forming minerals, dark minerals (B), quartz (Q) and feldspars with muscovite (F).
Grebennikov (2014) 5Fe <sub>2</sub> O <sub>3t</sub> - Na <sub>2</sub> O+K <sub>2</sub> O - Na <sub>2</sub> O+K <sub>2</sub> O - 5(CaO+MgO)	Grebennikov	Gr	5Fe <sub>2</sub> O <sub>3t</sub> – Na <sub>2</sub> O+K <sub>2</sub> O – 5(CaO+MgO) ternary	Plot of Grebennikov (2014) serves for classification of A-type granites and related felsic volcanic rocks (SiO <sub>2</sub> > 67 wt.%)

<sup>1</sup>Scope: Gr: granitoids, B: basaltoids, UB: ultrabasic rocks, U: (moreless) universal (appropriate for a range of compositions).

## REFERENCES

- Agrawal S, Guevara M, Verma S (2008). Tectonic discrimination of basic and ultrabasic volcanic rocks through log-transformed ratios of immobile trace elements. *International Geology Review* **50**, 1057–1079.
- Batchelor, R. A. & Bowden, P. (1985). Petrogenetic interpretation of granitoid rock series using multicationic parameters. *Chemical Geology* **48**, 43–55.
- Cabanis, B. & Lecolle, M. (1989). Le diagramme La/10-Y/15-Nb/8: un outil pour la discrimination des séries volcaniques et la mise en évidence des processus de mélange et/ou de contamination crustale. *Comptes rendus de l'Académie des sciences.Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre* **309**, 2023–2029.
- Cox, K. G., Bell, J. D. & Pankhurst, R. J. (1979). *The Interpretation of Igneous Rocks*. George Allen & Unwin.
- De La Roche, H., Leterrier, J., Grandclaude, P. & Marchal, M. (1980). A classification of volcanic and plutonic rocks using R<sub>1</sub>R<sub>2</sub>-diagram and major element analyses – its relationships with current nomenclature. *Chemical Geology* **29**, 183–210.
- Debon, F. & Le Fort, P. (1983). A chemical–mineralogical classification of common plutonic rocks and associations. *Transactions of the Royal Society of Edinburgh, Earth Sciences* **73**, 135–149.
- Debon, F. & Le Fort, P. (1988). A cationic classification of common plutonic rocks and their magmatic associations: principles, method, applications. *Bulletin de Minéralogie* **111**, 493–510.
- Enrique, P. (2018). Clasificación normativa de las rocas plutónicas saturadas y sobresaturadas en sílice basada en la clasificación modal QAP: el diagrama 2Q–(or + ab)–4an. *Geogaceta* **63**, 95–98.
- Frost, B.R. & Frost, C.D. (2008) A geochemical classification for feldspathic igneous rocks. *Journal of Petrology* **49**, 1955–1969.
- Frost, B. R., Barnes, C. G., Collins, W. J., Arculus, R. J., Ellis, D. J. & Frost, C. D. (2001). A geochemical classification for granitic rocks. *Journal of Petrology* **42**, 2033–2048.
- Grebennikov, A. V. (2014). A-type granites and related rocks: petrogenesis and classification. *Russian Geology and Geophysics* **55**, 1353–1366.
- Harris N B W, Pearce J A, Tindle A G (1986) Geochemical characteristics of collision- zone magmatism. In: Coward M P, Ries A C (eds) Collision Tectonics. Geological Society London Special Publications 19, pp 67-81.
- Hastie, A. R., Kerr, A. C., Pearce, J. A. & Mitchell, S. F. (2007). Classification of altered volcanic island arc rocks using immobile trace elements: development of the Th Co discrimination diagram. *Journal of Petrology* **48**, 2341–2357.
- Hollocher, K., Robinson, P., Walsh, E. & Roberts, D. (2012). Geochemistry of amphibolite-facies volcanics and gabbros of the Storen Nappe in extensions west and southwest of Trondheim, western gneiss region, Norway: A key to correlations and paleotectonic settings. *American Journal of Science* **312**, 357–416.
- Irvine, T. N. & Baragar, W. R. A. (1971). A guide to the chemical classification of the common volcanic rocks. *Canadian Journal of Earth Sciences* **8**, 523–548.
- Jensen, L. S. (1976). *A New Cation Plot for Classifying Subalkalic Volcanic Rocks*. Ontario Geological Survey Miscellaneous Paper **66**.
- Laurent, O., Martin, H., Moyen, J. F. & Doucelance, R. (2014). The diversity and evolution of late-Archean granitoids: evidence for the onset of 'modern-style' plate tectonics between 3.0 and 2.5 Ga. *Lithos* **205**, 208–235.
- Le Bas, M. J., Le Maitre, R. W., Streckeisen, A. & Zanettin, B. (1986). A chemical classification of volcanic rocks based on the total alkali–silica diagram. *Journal of Petrology* **27**, 745–750.
- Le Maitre, R. W. (1989). *A Classification of Igneous Rocks and Glossary of Terms. Recommendations of the IUGS Commission on the Systematics of Igneous Rocks*. Oxford: Blackwell.
- Maniar, P. D. & Piccoli, P. M. (1989). Tectonic discriminations of granitoids. *Geological Society of America Bulletin* **101**, 635–643.
- Martin, H. (1986) Effect of steeper Archean geothermal gradient on geochemistry of subduction-zone magmas. *Geology* **14**, 753–756.
- Meschede, M. (1986). A method of discriminating between different types of mid-ocean ridge basalts and continental tholeiites with the Nb–Zr–Y diagram. *Chemical Geology* **56**, 207–218.

- Middlemost, E. A. K. (1985). *Magmas and Magmatic Rocks*. London: Longman.
- Middlemost, E. A. K. (1985). Naming materials in the magma/igneous rock system. *Earth-Sciences Reviews* **37**, 215–224.
- Mielke, P. & Winkler, H. G. F. (1979). Eine bessere Berechnung der Mesonorm für granitische Gesteine. *Neues Jahrbuch für Mineralogie, Monatshefte* 471–480.
- Miyashiro A. (1974). Volcanic rock series in island arcs and active continental margins. *American Journal of Science* **274**, 321–355.
- Mullen, E. D. (1983). MnO/TiO<sub>2</sub>/P<sub>2</sub>O<sub>5</sub>: a minor element discriminant for basaltic rocks of oceanic environments and its implications for petrogenesis. *Earth and Planetary Science Letters* **62**, 53–62.
- Müller, D., Rock, N. M. S. & Groves, D. I. (1992). Geochemical discrimination between shoshonitic and potassic volcanic rocks in different tectonic settings: a pilot study. *Mineralogy and Petrology* **46**, 259–289.
- O'Connor, J. T. (1965). A classification for quartz-rich igneous rocks based on feldspar ratios. In: *US Geological Survey Professional Paper B525*. USGS, 79–84.
- Ohta, T. & Arai, H. (2007) Statistical empirical index of chemical weathering in igneous rocks: a new tool for evaluating the degree of weathering. *Chemical Geology* **240**, 280–297.
- Paulick, H., Bach, W., Godard, M., De Hoog, J. C. M., Suhr, G. & Harvey, J. (2006). Geochemistry of abyssal peridotites (Mid-Atlantic Ridge, 15°20'N, ODP Leg 209): Implications for fluid/rock interaction in slow spreading environments. *Chemical Geology* **234**, 179–210.
- Pearce, J. A. (1982). Trace element characteristics of lavas from destructive plate boundaries. In: Thorpe R.S. (ed.) *Andesites: Orogenic Andesites and Related Rocks*. John Wiley & Sons, Chichester, pp. 525–548, ISBN 0 471 28034 8
- Pearce, J. A. (1996). A user's guide to basalt discrimination diagrams. In: Wyman, D. A. (ed.) *Trace Element Geochemistry of Volcanic Rocks: Applications for Massive Sulphide Exploration*. Geological Association of Canada, Short Course Notes 12, 79–113.
- Pearce, J. A. (2008). Geochemical fingerprinting of oceanic basalts with applications to ophiolite classification and the search for Archean oceanic crust. *Lithos* **100**, 14–48.
- Pearce, J. A. & Cann, J. R. (1973). Tectonic setting of basic volcanic rocks determined using trace element analyses. *Earth and Planetary Science Letters* **19**, 290–300.
- Pearce, J. A. & Norry, M. J. (1979). Petrogenetic implications of Ti, Zr, Y, and Nb variations in volcanic rocks. *Contributions to Mineralogy and Petrology* **69**, 33–47.
- Pearce, T. H., Gorman, B. E. & Birkett, T. C. (1977). The relationship between major element geochemistry and tectonic environment of basic and intermediate volcanic rocks. *Earth and Planetary Science Letters* **36**, 121–132.
- Pearce, J. A., Harris, N. W. & Tindle, A. G. (1984). Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *Journal of Petrology* **25**, 956–983.
- Peccerillo, A. & Taylor, S. R. (1976). Geochemistry of Eocene calc-alkaline volcanic rocks from the Kastamonu area, Northern Turkey. *Contributions to Mineralogy and Petrology* **58**, 63–81.
- Ross, P. S. & Bédard, L. P. (2009). Magmatic affinity of modern and ancient subalkaline volcanic rocks determined from trace-element discriminant diagrams. *Canadian Journal of Earth Sciences* **46**, 823–839.
- Saccani, E. (2015). A new method of discriminating different types of post-Archean ophiolitic basalts and their tectonic significance using Th–Nb and Ce–Dy–Yb systematics. *Geoscience Frontiers* **6**, 481–501.
- Schandl, E. S. & Gorton, M. P. (2002). Application of high field strength elements to discriminate tectonic settings in VMS environments. *Economic Geology* **97**, 629–642.
- Shand, S. J. (1943). *Eruptive Rocks. Their Genesis, Composition, Classification, and Their Relation to Ore-Deposits with a Chapter on Meteorite*. New York: John Wiley & Sons.
- Shervais, J. W. (1982). Ti–V plots and the petrogenesis of modern and ophiolitic lavas. *Earth and Planetary Science Letters* **59**, 101–118.
- Streckeisen, A. (1974). Classification and nomenclature of plutonic rocks. *Geologische Rundschau* **63**, 773–786.
- Streckeisen, A. (1978). IUGS Subcommittee on the Systematics of Igneous Rocks: Classification and nomenclature of volcanic rocks, lamprophyres, carbonatites and melilitic rocks; recommendation and suggestions. *Neues Jahrbuch für Mineralogie, Abhandlungen* **134**, 1–14.
- Streckeisen, A. & Le Maitre, R. W. (1979). A chemical approximation to the modal QAPF classification of the igneous rocks. *Neues Jahrbuch für Mineralogie, Abhandlungen* **136**, 169–206.

- Sylvester P J (1989). Post-collisional alkaline granites. *Journal of Geology* **97**, 261–280.
- Verma S P, Guevara M, Agrawal S (2006). Discriminating four tectonic settings: Five new geochemical diagrams for basic and ultrabasic volcanic rocks based on log-ratio transformation of major-element data. *Journal of Earth System Science* **115**, 485–528.
- Villaseca, C., Barbero, L. & Herreros, V. (1998). A re-examination of the typology of peraluminous granite types in intracontinental orogenic belts. *Transactions of the Royal Society of Edinburgh, Earth Sciences* **89**, 113–119.
- Whalen J, B., Currie K. L. & Chappell B. W. (1987). A-type granites: geochemical characteristics, discrimination and petrogenesis. *Contributions to Mineralogy and Petrology* **95**, 407–419.
- Wilson, M. (1989). *Igneous Petrogenesis*. London: Unwin Hyman.
- Winchester, J. A. & Floyd, P. A. (1977). Geochemical discrimination of different magma series and their differentiation products using immobile elements. *Chemical Geology* **20**, 325–343.
- Wood, D. A. (1980). The application of a Th–Hf–Ta diagram to problems of tectonomagmatic classification and to establishing the nature of crustal contamination of basaltic lavas of the British Tertiary volcanic province. *Earth and Planetary Science Letters* **50**, 11–30.